

Definitions: Let $\Gamma = (V, E)$ be an undirected graph and let u and v be (not necessarily distinct) vertices in Γ .

- A *path of length n* from u to v in Γ is a sequence $\langle v_0, e_1, v_1, e_2, \dots, v_{n-2}, e_{n-1}, v_{n-1}, e_n, v_n \rangle$ with $v_0 = u$, $v_n = v$, and each edge e_i for $1 \leq i \leq n$ connects v_{i-1} and v_i .
- Given such a path, the sequence $\langle v_0, v_1, \dots, v_{n-1}, v_n \rangle$ is the *vertex sequence* of the path, and the sequence $\langle e_1, e_2, \dots, e_{n-1}, e_n \rangle$ is the *edge sequence* of the path.
- A path is said to *pass through* the vertices v_1, v_2, \dots, v_{n-1} . v_0 and v_n are called the *endpoints* of the path. A path is said to *traverse* its edges e_1, e_2, \dots, e_n .
- In a simple graph, we may use the vertex sequence for a path to identify the path, but in a multigraph, the vertex sequence may not give enough information to completely specify the path.
- A path is a *circuit* if $v_0 = v_n$ (that is, if it begins and ends at the same vertex) and the length of the path is greater than zero.
- A path (or circuit) is *simple* if it does not contain the same edge more than once.
- We say that Γ is *connected* if there is at least one path between every (distinct) pair of vertices in V .
- A *connected component* of a graph Γ is a proper subgraph of Γ that is connected and which is not properly contained in any other connected subgraph of Γ . That is, it is a maximal connected proper subgraph of Γ .

Definitions: Let $\Gamma = (V, E)$ be a directed graph and let u and v be (not necessarily distinct) vertices in Γ .

- A *path of length n* from u to v in Γ is a sequence $\langle v_0, e_1, v_1, e_2, \dots, v_{n-2}, e_{n-1}, v_{n-1}, e_n, v_n \rangle$ with $v_0 = u$, $v_n = v$, and each edge e_i for $1 \leq i \leq n$ connects v_{i-1} to v_i (as a directed edge).
- A (directed) path is a *circuit* if $v_0 = v_n$ (that is, if it begins and ends at the same vertex) and the length of the path is greater than zero.
- A path (or circuit) is *simple* if it does not contain the same edge more than once.
- A directed graph Γ is *strongly connected* if for every pair of vertices u, v in Γ , there is a path from u to v and there is also a path from v to u .
- A directed graph Γ is *weakly connected* if there its underlying undirected graph is connected.

Theorem: There is a simple path between every pair of distinct vertices in a connected undirected graph.

Proof: Let u and v be distinct vertices in a connected undirected graph Γ . By the definition of connected, there is at least on path from u to v . Let $\langle v_0, v_1, \dots, v_{n-1}, v_n \rangle$ be the vertex sequence of a path from u to v of shortest possible length.

Claim: this shortest path is simple.

To see this, suppose that it is not simple. Then $v_i = v_j$ for some pair of vertices in the vertex sequence with $0 \leq i < j$ (this is because whenever there is a repeated edge, at least one of the vertices along that edge must be visited twice). But then we may remove a segment of the path to get a path of shorter length with vertex sequence $v_0, v_1, \dots, v_{i-1}, v_j, v_{j+1}, \dots, v_n$. If this were the case, we would have found a path from u to v that is shorter than our original path, contradicting our original assumption. \square

Definitions:

- An *Euler path* in a graph Γ is a simple path containing every edge of Γ . An *Euler circuit* is an Euler path that is a circuit (i.e that begins and ends at the same vertex).
- A *Hamiltonian path* in a graph Γ is a simple path that passes through every vertex of Γ exactly once. A *Hamiltonian circuit* is a Hamiltonian path that is a circuit (i.e that begins and ends at the same vertex).