

**Definition:** A function  $f$  is **continuous** at an interior point  $c$  of its domain if:


- $f(c)$  is defined
- $\lim_{x \rightarrow c} f(x)$  exists
- $\lim_{x \rightarrow c} f(x) = f(c)$

A function  $f$  is **continuous** at a left endpoint  $a$  or a right endpoint  $b$  of its domain if:

- $f(a)$  is defined,  $\lim_{x \rightarrow a^+} f(x)$  exists, and  $\lim_{x \rightarrow a^+} f(x) = f(a)$ .  
or
- $f(b)$  is defined,  $\lim_{x \rightarrow b^-} f(x)$  exists, and  $\lim_{x \rightarrow b^-} f(x) = f(b)$ .

**Notes:**


- We really only need to check the third condition, since the first two are implied by the third.
- If one (or more) of the three conditions in the definition above are not satisfied, we say that  $f$  is **discontinuous** at the corresponding value or that  $f$  has a **discontinuity** at that value.
- We can extend the definition above to say that a function  $f(x)$  is continuous on an open interval  $(a, b)$  if it is continuous at every interior point of this interval. We can further extend it to say that a function  $f(x)$  is continuous on a closed interval  $[a, b]$  if it is continuous at every interior point and is continuous at both endpoints as well.



We often classify discontinuities as one of four main types.

1. If  $\lim_{x \rightarrow c} f(x)$  exists but  $f(c)$  is undefined or attains a different value, we say that  $f$  has a **removable discontinuity** at  $c$ . This is due to the fact that redefining the value of  $f(c)$  would make  $f$  continuous at  $c$ , thereby “removing” the discontinuity at this input value. Take a moment to identify the input values that exhibit this type of discontinuity in the figure above.
2. If  $\lim_{x \rightarrow c^-} f(x) \neq \lim_{x \rightarrow c^+} f(x)$ , we say that  $f$  has a **jump discontinuity** at  $c$ . Take a moment to identify the input values that exhibit this type of discontinuity in the figure above.
3. If either  $\lim_{x \rightarrow c^-} f(x)$  or  $\lim_{x \rightarrow c^+} f(x)$  attains an infinite value ( $\infty$  or  $-\infty$ ), then we say that  $f$  has an **infinite discontinuity** at  $c$ . Take a moment to identify the input values that exhibit this type of discontinuity in the figure above.

4. The fourth type of discontinuity that can occur is a bit harder to explain. It is often called discontinuity due to **oscillation**. To see an example of this type of discontinuity, consider the graph of the function  $f(x) = \sin\left(\frac{1}{x}\right)$  at  $x = 0$ .



**Useful Facts:**

- A polynomial function  $f(x)$  is continuous at every real number  $c$ .
- A rational function  $q(x) = \frac{f(x)}{g(x)}$  where  $f$  and  $g$  are polynomials is continuous at every number  $c$  for which  $g(c) \neq 0$ .

**Theorem 8:** If  $f(x)$  and  $g(x)$  are continuous at  $x = c$ , then the following are also continuous at  $x = c$ :

- the sum  $(f + g)(x)$
- the difference  $(f - g)(x)$
- $k \cdot f(x)$  for any constant  $k$ .
- $\sqrt[n]{f(x)}$  for any positive integer  $n$  (assuming  $f(x)$  is positive on an interval containing  $c$  when  $n$  is even).
- the product  $(fg)(x)$
- the quotient  $\left(\frac{f}{g}\right)(x)$ , provided  $g(c) \neq 0$
- $[f(x)]^n$  for any positive integer  $n$ .

**Theorem 10:** If  $\lim_{x \rightarrow c} g(x) = b$  and  $f$  is continuous at  $b$ , then  $\lim_{x \rightarrow c} f(g(x)) = f(b) = f\left(\lim_{x \rightarrow c} g(x)\right)$

**Theorem 9:** If  $g$  is continuous at  $c$  and if  $f$  is continuous at  $b = g(c)$ , then

- $\lim_{x \rightarrow c} f(g(x)) = f\left(\lim_{x \rightarrow c} g(x)\right) = f(g(c))$
- the composite function  $f \circ g$  is continuous at  $c$ .

**The Intermediate Value Theorem:** If  $f$  is continuous on a closed interval  $[a, b]$  and if  $w$  is any number between  $f(a)$  and  $f(b)$ , then there is at least one number  $c$  in  $[a, b]$  such that  $f(c) = w$ .

**Example:**