

1. Suppose you throw a ball vertically upward. If you release the ball 7 feet above the ground at an initial speed of 48 feet per second, how high will the ball travel? (Assume gravity is $-32ft/sec^2$)

We know that $a(t) = -32$, $v(0) = 48$, and $s(0) = 7$.

Antidifferentiating, $v(t) = -32t + C$, so $v(0) = 48 = -32(0) + C$, so $C = 48$, and $v(t) = -32t + 48$.

Antidifferentiating again, $s(t) = -16t^2 + 48t + D$, so $s(0) = 7 = D$, and $s(t) = -16t^2 + 48t + 7$.

The max height occurs when $v(t) = 0$, that is, when $-32t + 48 = 0$, or $32t = 48$, so when $t = \frac{48}{32} = \frac{3}{2}$.
[Notice $s''(t) = a(t) < 0$, so we know it is a maximum]

Thus, the max height is: $s(\frac{3}{2}) = -16(\frac{3}{2})^2 + 48(\frac{3}{2}) + 7 = 43$, or 43 feet.

2. Find each of the following indefinite integrals:

$$(a) \int \frac{x^{\frac{3}{2}} - 7x^{\frac{1}{2}} + 3}{x^{\frac{1}{2}}} dx$$

$$= \int x - 7 + 3x^{-\frac{1}{2}} dx = \frac{1}{2}x^2 - 7x + 6x^{\frac{1}{2}} + C$$

$$(b) \int \sin^3 x \cos x dx$$

Let $u = \sin x$. Then $du = \cos x dx$, and we have $\int u^3 du = \frac{1}{4}u^4 + C$.

Thus the indefinite integral is: $\frac{1}{4}\sin^4 x + C$.

$$(c) \int 5x(x^2 + 1)^8 dx$$

Let $u = x^2 + 1$. Then $du = 2x dx$, or $\frac{1}{2} du = dx$, so we have $\int \frac{5}{2}u^8 du = \frac{5}{2} \frac{1}{9}u^9 + C = \frac{5}{18}u^9$.

Thus the indefinite integral is: $\frac{5}{18}(x^2 + 1)^9 + C$.

$$(d) \int \frac{x}{\sqrt{x+1}} dx$$

This is a trickier substitution problem. Let $u = x + 1$. Then $u - 1 = x$, and $du = dx$.

Thus we have the indefinite integral: $\int \frac{u-1}{\sqrt{u}} du = \int u^{\frac{1}{2}} - u^{-\frac{1}{2}} du = \frac{2}{3}u^{\frac{3}{2}} - 2u^{\frac{1}{2}} + C$.

Thus the indefinite integral is: $\frac{2}{3}(x+1)^{\frac{3}{2}} - 2(x+1)^{\frac{1}{2}} + C$.

3. Solve the following initial value problems under the given initial conditions:

$$(a) \frac{dy}{dx} = \sin x + x^2; y = 5 \text{ when } x = 0$$

Antidifferentiating, $y = -\cos x + \frac{1}{3}x^3 + C$.

Therefore, $5 = -\cos(0) + \frac{1}{3}(0)^3 + C$, or $5 = -1 + 0 + C$, so $C = 6$.

Therefore, $y = -\cos x + \frac{1}{3}x^3 + 6$.

(b) $g''(x) = 4 \sin(2x) - \cos(x); g'(\frac{\pi}{2}) = 3; g(\frac{\pi}{2}) = 6$

Antidifferentiating, $g'(x) = -2 \cos(2x) - \sin x + C$

Therefore, $3 = -2 \cos(\pi) - \sin \frac{\pi}{2} + C = -2(-1) - 1 + C$, so $3 = 2 - 1 + C$, or $C = 2$.

Hence $g'(x) = -2 \cos(2x) - \sin x + 2$. But then $g(x) = -\sin(2x) + \cos x + 2x + D$.

Moreover, $6 = -\sin(\pi) + \cos \frac{\pi}{2} + 2(\frac{\pi}{2}) + D$, or $6 = 0 + 0 + \pi + D$, so $D = 6 - \pi$

Thus $g(x) = -\sin(2x) + \cos x + 2x + 6 - \pi$.

4. Express the following in summation notation:

(a) $2 + 5 + 10 + 17 + 26 + 37 = \sum_{k=1}^6 (k^2 + 1)$

(b) $x^2 + \frac{x^3}{4} + \frac{x^4}{9} + \dots + \frac{x^{11}}{100} = \sum_{k=1}^{10} \frac{x^{k+1}}{k^2}$

5. Evaluate the following sums:

(a) $\sum_{k=2}^5 k^2(k+1)$
 $= \sum_{k=2}^5 k^3 + k^2 = (8 + 4) + (27 + 9) + (64 + 16) + (125 + 25) = 278$

(b) $\sum_{k=3}^{20} k^3 - k^2$
 $= \sum_{k=1}^{20} k^3 - \sum_{k=1}^{20} k^2 - \sum_{k=1}^2 k^3 + \sum_{k=1}^2 k^2$
 $= \left(\frac{(20)(21)}{2}\right)^2 - \frac{(20)(21)(41)}{6} - 1 - 8 + 1 + 4 = (210)^2 - 2870 - 9 + 5 = 41,226$

6. Express the following sums in terms of n :

(a) $\sum_{k=1}^n 3k^2 - 2k + 10$
 $= 3 \sum_{k=1}^n k^2 - 2 \sum_{k=1}^n k + \sum_{k=1}^n 10 = 3 \frac{(n)(n+1)(2n+1)}{6} - 2 \frac{(n)(n+1)}{2} + 10n$
 $= n^3 + \frac{3n^2}{2} + \frac{n}{2} - n^2 - n + 10n = n^3 + \frac{n^2}{2} + \frac{19n}{2}$

(b) $\sum_{k=3}^n k(k^2 - 1)$
 $= \sum_{k=1}^n k^3 - \sum_{k=1}^n k - \left(\sum_{k=1}^2 k^3 - \sum_{k=1}^2 k \right)$
 $= \left(\frac{(n)(n+1)}{2}\right)^2 - \frac{(n)(n+1)}{2} - (1 + 8 - 1 - 2) = \frac{n^4 + 2n^3 + n^2}{4} - \frac{n^2 + n}{2} - 6 = \frac{n^4}{4} + \frac{n^3}{2} - \frac{n^2}{4} - \frac{n}{2} - 6.$

7. Consider $f(x) = 3x^2 - 5$ in the interval $[3, 7]$

- (a) Find a summation formula that gives an estimate the definite integral of f on $[3, 7]$ using n equal width rectangles and using midpoints to give the height of each rectangle. You do not have to evaluate the sum or find the exact area.

Notice that $\Delta x = \frac{7-3}{n} = \frac{4}{n}$. Since we want to use midpoints for our heights, $x_k = 3 + k\Delta x - \frac{\Delta x}{2} = 3 + \frac{4k-2}{n}$.

$$\text{Therefore, } A_n = \sum_{k=1}^n f(x_k)\Delta x = \sum_{k=1}^n \left[3 \left(3 + \frac{4k-2}{n} \right)^2 - 5 \right] \left(\frac{4}{n} \right)$$

- (b) Find the norm of the partition $P : 3 < 3.5 < 5 < 6 < 6.25 < 7$

The norm is the widest gap in the partition: $5 - 3.5 = 1.5$

- (c) Find the approximation of the definite integral of f on $[3, 7]$ using the Riemann sum for the partition P given in part (b).

$$A \approx \sum_{k=1}^5 f(x_k)\Delta x_k = f(3.25)(.5) + f(4.25)(1.5) + f(5.5)(1) + f(6.125)(.25) + f(6.625)(.75) = (26.6875)(.5) + (49.1875)(1.5) + (85.75)(1) + (107.546875)(.25) + (126.671875)(.75) = 294.765625$$

8. Assume f is continuous on $[-5, 3]$, $\int_{-5}^{-1} f(x) dx = -7$, $\int_{-1}^3 f(x) dx = 4$, and $\int_1^3 f(x) dx = 2$. Find:

(a) $\int_3^{-1} f(x) dx = - \int_{-1}^3 f(x) dx = -4$

(b) $\int_{-5}^1 f(x) dx = \int_{-5}^{-1} f(x) dx + \int_{-1}^3 f(x) dx - \int_1^3 f(x) dx = -7 + 4 - 2 = -5$

(c) $\int_{-5}^3 f(x) dx = \int_{-5}^{-1} f(x) dx + \int_{-1}^3 f(x) dx = -7 + 4 = -3$

(d) $\int_{-1}^{-1} f(x) dx = 0$

- (e) Find the average value of f on $[-5, -1]$

$$= \frac{1}{-1 - (-5)} \int_{-5}^{-1} f(x) dx = \frac{1}{4} \cdot (-7) = -\frac{7}{4}$$

9. Evaluate the following:

(a) $\int_1^4 x^3 + \frac{1}{\sqrt{x}} + 2 dx$
 $= \frac{1}{4}x^4 + 2x^{\frac{1}{2}} + 2x \Big|_1^4 = \left[\frac{1}{4}4^4 + 2 \cdot 4^{\frac{1}{2}} + 2(4) \right] - \left[\frac{1}{4}1^4 + 2 \cdot 1^{\frac{1}{2}} + 2(1) \right] = 76 - 4.25 = 71.75$

(b) $\int_0^1 x^2(2x^3 + 1)^2 dx$

Let $u = 2x^3 + 1$. Then $du = 6x^2$, or $\frac{1}{6}du = dx$.

Notice $2(0)^3 + 1 = 1$, and $2(1)^3 + 1 = 3$

$$\text{Then we have } \int_1^3 \frac{1}{6}u^2 du = \frac{1}{18}u^3 \Big|_1^3 = \frac{1}{18}(3^3 - 1^3) = \frac{26}{18} = \frac{13}{9}.$$

$$(c) \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \sin^3(2x) \cos(2x) dx$$

Let $u = \sin(2x)$. Then $du = 2 \cos(2x) dx$ or $\frac{1}{2} du = \cos(2x) dx$

Notice that $\sin(2 \cdot \frac{\pi}{6}) = \sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$, while $\sin \pi = 0$

$$\text{Then we have } \int_{\frac{\sqrt{3}}{2}}^0 \frac{1}{2} u^3 du = \frac{1}{4} u^4 \Big|_{\frac{\sqrt{3}}{2}}^0 = 0 - \frac{1}{8} \cdot \left(\frac{\sqrt{3}}{2}\right)^4 = -\frac{9}{128}$$

$$(d) \int_{-\pi}^{\pi} \sin x dx = 0, \text{ since } \sin x \text{ is an odd function.}$$

$$(e) \frac{d}{dx} \left(\int_1^3 t \sqrt{t^2 - 1} dt \right) = 0, \text{ since a definite integral gives a constant, and the derivative of a constant is zero.}$$

$$(f) \int_1^3 \left[\frac{d}{dt} \left(t \sqrt{t^2 - 1} \right) dt \right] = (3\sqrt{3^2 - 1}) - (1\sqrt{1^2 - 1}) = 3\sqrt{8} = 6\sqrt{2}$$

$$10. \text{ Suppose } G(x) = \int_2^x \frac{1}{t^2 + 1} dt$$

$$(a) \text{ Find } G'(2) = \frac{1}{2^2 + 1} = \frac{1}{5}$$


$$(b) \text{ Find } G'(x^2) = \frac{1}{x^4 + 1} \text{ [Note: } G'(x^2) \neq \frac{d}{dx} G(x^2)\text{]}$$

$$(c) \text{ Find } G''(3)$$

Notice that $\frac{d}{dx} \frac{1}{x^2 + 1} = \frac{-2x}{(x^2 + 1)^2}$.

$$\text{Thus } G''(3) = \frac{-2(3)}{(3^2 + 1)^2} = \frac{-6}{(10)^2} = \frac{-3}{50}.$$

$$11. \text{ Given the following graph of } f(x) \text{ and the fact that } G(x) = \int_{-2}^x f(t) dt:$$



$$(a) \text{ Find } G(6) = \int_{-2}^6 f(t) dt = 2.5 + 3.5 + \frac{1}{2}(2)(4) - \frac{1}{2}(4)(2) = 6 \text{ [Compute area directly]}$$

$$(b) \text{ Find } G'(6) = f(6) = 0$$

$$(c) \text{ Find } G''(6) = f'(6) = 1 \text{ [Compute slope of line segment on graph at } x = 6\text{]}$$

12. Find the area between the curves:

(a) $y = x^2 + 1$ and $y = 3x - 1$

First notice that these curves intersect when $x^2 + 1 = 3x - 1$, or when $x^2 - 3x + 2 = 0$. That is, when $(x - 2)(x - 1) = 0$, or when $x = 2$ and $x = 1$.

Next, notice that $3x - 1 \geq x^2 + 1$ on $[1, 2]$. Thus the area between these curves is given by:

$$\int_1^2 (3x - 1) - (x^2 + 1) dx = \int_1^2 -x^2 + 3x - 2 dx = -\frac{x^3}{3} + \frac{3}{2}x^2 - 2x \Big|_1^2 = \frac{1}{6}$$

(b) $y = x^2 - 1$ and $y = 1 - x$ on $[0, 2]$

First notice that these curves intersect when $x^2 - 1 = 1 - x$, or when $x^2 + x - 2 = 0$. That is, when $(x + 2)(x - 1) = 0$, or when $x = -2$ and $x = 1$.

Next, notice that $1 - x \geq x^2 - 1$ on $[0, 1]$ while $x^2 - 1 \geq 1 - x$ on $[1, 2]$. Thus the area between these curves is given by:

$$\begin{aligned} \int_0^1 (1 - x) - (x^2 - 1) dx + \int_1^2 (x^2 - 1) - (1 - x) dx &= \int_0^1 -x^2 - x + 2 dx + \int_1^2 x^2 + x - 2 dx \\ &= \left(-\frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x \Big|_0^1 \right) + \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 - 2x \Big|_1^2 \right) = 3. \end{aligned}$$

(c) $y = x$, $y = 2$, $y + x = 6$, and $y = 0$

This integral is a bit easier if we slice the area horizontally and integrate with respect to y , so, solving for x in terms of y , we have $x = y$ and $x = 6 - y$, with $6 - y \geq y$ on $[0, 2]$. Therefore, the area of this region is given by:

$$\int_0^2 (6 - y) - (y) dy = \int_0^2 6 - 2y dy = 6y - y^2 \Big|_0^2 = 12 - 4 = 8$$

(d) $x = y^2$, $x = 4$

We will find the area by integrating with respect to y . Notice that $y^2 = 4$ when $y = \pm 2$. Also, $4 \geq y^2$ on $[-2, 2]$. Therefore, the area of this region is given by:

$$\int_{-2}^2 4 - y^2 dy = 4y - \frac{1}{3}y^3 \Big|_{-2}^2 = \frac{32}{3}.$$

13. (a) Use the Trapezoidal Rule with $n = 4$ to approximate $\int_0^4 2x^3 dx$

$$A \approx \frac{4-0}{4 \cdot 2} [f(0) + 2f(1) + 2f(2) + 2f(3) + f(4)] = \frac{1}{2} [0 + 2(2 \cdot 1) + 2(2 \cdot 8) + 2(2 \cdot 27) + 2 \cdot 64] = \frac{1}{2} [272] = 136$$

(b) Use the Fundamental Theorem of Calculus to find $\int_0^4 2x^3 dx$ exactly. How far off was your estimate?

$$\int_0^4 2x^3 dx = \frac{1}{2} x^4 \Big|_0^4 = \frac{1}{2} 4^4 - 0 = 128$$

The error in our approximation is $136 - 128 = 8$