- 1. Find the derivative of each. Are the solutions what you expected? If not, explain how and why.
 - (a) $f(x) = (4x^3 + 3x 5) \sin x$ (b) $p(t) = \frac{5 - 3t^2}{t^3 + 4t - 1} \cos^2 t$ (c) $g(\theta) = \theta \cot(\theta^2 + 3\theta - 1)$
- 2. Use the functions from problem 1 to find the instantaneous rate of change for the function at the given value. (Express the solutions in a *reasonable* form.)
 - (a) f(x) when x = 1
 - (b) p(t) when $t = \pi/6$
 - (c) $g(\theta)$ when $\theta = 0.41$

3. Given $f(x) = \frac{x^4}{4} - 3x^2 - 2x + 1$.

- (a) Graph f and its first, second, third, and fourth derivatives on the same coordinate plane. Show an appropriately labeled legend.
- (b) What is true about the function f when the first derivative is negative?
- (c) What is true about the first derivative when the second derivative is negative?
- (d) How does the relationships between the graphs of each succeeding derivative illustrate the expected result from the Power Rule?

4. Given
$$g(x) = (x^2 - x - 6) \sin\left(\frac{x^2}{10}\right)$$
 on $[-\pi, \pi]$.

- (a) Determine the coordinates of all points where g has horizontal tangents in the given interval.
- (b) Determine the coordinates of all points in the given interval where the slope of a tangent line to the graph of g is -4.

5. Given
$$R(t) = \frac{5 - 3t + 4t^3 - t^4}{t^2 - t + 2}$$
.

- (a) Determine the slope of the tangent line to the graph of R when t = 1, 2.1, and 4.31.
- (b) Determine the interval(s) when the second derivative is positive.