

Math 229 Functions 06/02/2008

#### Recall:

More formally, a function f from a domain set D to a set E is a correspondence that assigns to each element x of D exactly one element y of E. We call x the **argument** of f and y the **value** of f at x. The **range** of f is the subset R of E consisting of all y values that corresponding to an x in the domain D.





- $\bullet$  To evaluate a function, we input an x-value and find the corresponding value by applying the "rule" for the function to that input.
- $\bullet$  Sometimes we also want to work backwards, that is, given an **output**, we try to find the input(s) that lead to that particular output.
- To find the domain of a function, we carefully analyze the function "rule" and find any x values that do not have corresponding outputs. Two things we look for in particular are division by zero and even roots of negative numbers.

## Example 1:

Suppose  $f(x) = \frac{x+1}{x-1}$ . Then:

$$f(2) = \frac{2+1}{2-1} = \frac{3}{1} = 3$$

$$f(-1) = \frac{-1+1}{-1-1} = \frac{0}{-2} = 0$$

$$f(2a-1) = \frac{2a-1+1}{2a-1-1} = \frac{2a}{2a-2} = \frac{a}{a-1}$$

If f(x) = 2, that what is x?

$$\frac{x+1}{x-1} = 2$$
, so  $x + 1 = 2(x - 1) = 2x - 2$ .

Then x + 3 = 2x, or 3 = x. Check:  $f(3) = \frac{3+1}{3-1} = \frac{4}{2} = 2$ .

The domain of f is ?

### Example 2:

Let  $g(x) = \frac{\sqrt{3x-3}}{x^2+2x-3}$  Find:

- $\bullet$  g(4)
- $\bullet$  g(1)
- the domain of g(x)

# II. Graphs of Functions

#### **Definition:**

The **graph** of a function is the set of all points (x, f(x)) (where x is in the domain D of f).

## The Vertical Line Test:

A graph of points in the plane is the graph of a function if and only if every vertical line intersects the graph at most once.


## **Definitions:**

A function is **increasing** on an interval I if  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$  in I.

A function is **decreasing** on an interval I if  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$  in I.

A function is **constant** on an interval I if  $f(x_1) = f(x_2)$  for all  $x_1, x_2$  in I.

# Example:



Find:

(a) f(4)

(b) 
$$x \text{ if } f(x) = 4$$

(c) the domain of f

(d) the range of f

(e) the intervals where f(x) is increasing