

Recall: The **Logarithm of x to the base b** is defined as follows: $y = \log_b x$ if and only if $x = b^y$. for $x > 0$ and $b > 0, b \neq 1$. A logarithm basically asks: "what power would I need to raise the base b to in order to get x as the result?"

Properties of logarithms: Let m and n be positive real numbers.

- | | |
|---|-----------------------|
| 1. $\log_b mn = \log_b m + \log_b n$ | 5. $\log_b b = 1$ |
| 2. $\log_b \frac{m}{n} = \log_b m - \log_b n$ | 6. $\log_b b^x = x$ |
| 3. $\log_b m^n = n \cdot \log_b m$ | 7. $b^{\log_b x} = x$ |
| 4. $\log_b 1 = 0$ | |

Examples: Use the Properties of Logarithms to expand the following:

- $\log_b 16 = \log_b 2^4 = 4 \log_b 2$
- $\log_b \frac{7}{16} = \log_b 7 - \log_b 16 = \log_b 7 - \log_b 2^4 = \log_b 7 - 4 \log_b 2$
- $$\log_b \left(\frac{(x+4)^3(x-1)^2}{\sqrt{x+1}} \right)$$

$$= \log_b ((x+4)^3(x-1)^2) - \log_b (\sqrt{x+1})$$

$$= \log_b (x+4)^3 + \log_b (x-1)^2 - \log_b (x+1)^{\frac{1}{2}}$$

$$= 3 \log_b (x+4) + 2 \log_b (x-1) - \frac{1}{2} \log_b (x+1)$$

Example: Use the Properties of Logarithms to combine the following into a single logarithm:

$$\begin{aligned} &= \frac{5}{2} \log_b (2x-7) - \log_b (3x+1) - \frac{3}{2} \log_b (x+1) \\ &= \frac{5}{2} \log_b (2x-7) - [\log_b (3x+1) + \frac{3}{2} \log_b (x+1)] \\ &= \log_b (2x-7)^{\frac{5}{2}} - [\log_b (3x+1) + \log_b (x+1)^{\frac{3}{2}}] \\ &= \log_b (2x-7)^{\frac{5}{2}} - [\log_b (3x+1)(x+1)^{\frac{3}{2}}] \\ &= \log_b \left(\frac{(2x-7)^{\frac{5}{2}}}{\log_b (3x+1) + \log_b (x+1)^{\frac{3}{2}}} \right) \end{aligned}$$

Examples: Solving Logarithmic Equations:

$$1. \log_3(x+6) - \log_3(x-2) = 2$$

$$\text{Then } \log_3 \left(\frac{x+6}{x-2} \right) = 2, \text{ so } 3^2 = \frac{x+6}{x-2}$$

$$\text{Therefore, } 9(x-2) = x+6, \text{ or } 9x-18 = x+6. \text{ Hence } 8x = 24, \text{ or } x = 3$$

$$\text{Check: } \log_3(3+6) - \log_3(3-2) = \log_3(9) - \log_3(1) = 2 - 0 = 2$$

$$2. \ln x = 1 - \ln(3x-2) - \ln e$$

$$\text{Then } \ln x + \ln(3x-2) = 1 - 1, \text{ or } \ln(x(3x-2)) = 0$$

$$\text{But then, exponentiating both sides: } e^{\ln(x(3x-2))} = e^0, \text{ or } x(3x-2) = 1$$

$$\text{Thus } 3x^2 - 2x - 1 = 0, \text{ or } (3x+1)(x-1) = 0.$$

$$\text{Hence } 3x = -1, \text{ or } x = -\frac{1}{3} \text{ and } x = 1$$

Notice that $x = -\frac{1}{3}$ does not check while $x = 1$ does check.