

1. Given the points  $A : (4, -2)$  and  $B : (-2, 7)$ :

(a) Find an equation for the line containing  $A$  and  $B$

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - (-2)}{-2 - 4} = \frac{9}{-6} = -\frac{3}{2}.$$

Using the point  $(-2, 7)$  and the point-slope formula,  $y - (7) = -\frac{3}{2}(x + 2)$ , or  $y - 7 = -\frac{3}{2}x - 3$ .

Therefore,  $y = -\frac{3}{2}x + 4$

(b) Find the line that is perpendicular to the line you found in part (a) and containing the point  $(1, -1)$

Since the slope of the previous line is  $m_1 = -\frac{3}{2}$ , a line that is perpendicular to the previous line has slope equal to the negative reciprocal  $m_2 = -\frac{1}{m_1} = \frac{2}{3}$ .

Also, since the line passes through  $(1, -1)$ , the equation of the line is given by:

$$y + 1 = \frac{2}{3}(x - 1) = \frac{2}{3}x - \frac{2}{3}$$


Thus,  $y = \frac{2}{3}x - \frac{5}{3}$

2. Find solutions to the inequality:  $\frac{x^2 - 1}{x^2 + x - 6} \leq 0$ .

Factoring, we have:  $\frac{(x + 1)(x - 1)}{(x + 3)(x - 2)} \leq 0$ .

Notice that the numerator is zero when  $x = 1$  or  $x = -1$ , and the denominator is zero when  $x = -3$  or  $x = 2$ .

Therefore, using sign analysis, we have the following sign diagram:



Thus the solution to this inequality, in interval notation, is:  $(-3, -1] \cup [1, 2)$

3. Given the function  $f(x) = \frac{1}{x - 2}$

(a) What is the domain of  $f$ ? Give your answer in interval notation.

Notice that  $f(x)$  is defined except when  $x = 2$ . Therefore, the domain, in interval notation, is:  $(-\infty, 2) \cup (2, \infty)$

(b) Find  $f(5)$  and  $f(2a + 4)$

$$f(5) = \frac{1}{5-2} = \frac{1}{3}. \text{ Similarly, } f(2a + 4) = \frac{1}{(2a+4)-2} = \frac{1}{2a+2} = \frac{1}{2(a+1)}$$

(c) Find  $\frac{f(a + h) - f(a)}{h}$  (be sure to simplify your answer).

$$\begin{aligned} \frac{f(a + h) - f(a)}{h} &= \frac{\frac{1}{a+h-2} - \frac{1}{a-2}}{h} = \frac{\frac{a-2}{a+h-2} - \frac{a+h-2}{a-2}}{h} = \frac{(a-2) - (a+h-2)}{(a+h-2)(a-2)} \cdot \frac{1}{h} = \frac{a-2-a-h+2}{(a+h-2)(a-2)} \cdot \frac{1}{h} = \\ &= \frac{-h}{(a+h-2)(a-2)} \cdot \frac{1}{h} = \frac{-1}{(a+h-2)(a-2)} \end{aligned}$$

4. Given that  $f(x) = \frac{1}{2x-3}$  and  $g(x) = \sqrt{x^2-9}$

(a) Find  $f \circ g(2)$

$f \circ g(2) = f(g(2)) = f(\sqrt{4-9}) = f(\sqrt{-5})$ , which is undefined.

(b) Find the domain of  $\frac{g}{f}$ ? Give your answer in interval notation.

To be in the domain of  $\frac{g}{f}$ , an  $x$ -value must be in the domain of both  $f(x)$  and  $g(x)$ , and we must also have  $g(x) \neq 0$ .

The domain of  $f(x)$  is all  $x$  except when  $2x-3=0$ , or  $2x=3$ . Thus, the domain is  $x \neq \frac{3}{2}$

The domain of  $g(x)$  is all  $x$  for which  $x^2-9 \geq 0$ , or  $x^2 \geq 9$ . Thus, we need  $|x| \geq 3$ . Hence  $x \geq 3$  or  $x \leq -3$ .

Finally, we need  $g(x) \neq 0$ , so  $x^2 \neq 9$ , or  $x \neq 3$  and  $x \neq -3$

Combining these, the domain is:  $(-\infty, -3) \cup (3, \infty)$

5. Find the exact value of each of the following:

(a)  $\sin\left(\frac{7\pi}{4}\right) = -\frac{\sqrt{2}}{2}$

(b)  $\cos\left(\frac{4\pi}{3}\right) = -\frac{1}{2}$

(c)  $\sin^{-1}\left(\frac{1}{\sqrt{2}}\right) = \frac{\pi}{4}$

(d)  $\cos^{-1}(-1) = \pi$

(e)  $\sin^{-1}\left(-\frac{\sqrt{3}}{2}\right) = -\frac{\pi}{3}$

6. Find all solutions to the following equations. Give the exact answers.

(a)  $2 \sin 3x = \sqrt{3}$

$\sin 3x = \frac{\sqrt{3}}{2}$ , so either  $3x = \frac{\pi}{3} + 2\pi k$  or  $3x = \frac{2\pi}{3} + 2\pi k$


Hence  $x = \frac{\pi}{9} + \frac{2\pi}{3}k$  or  $x = \frac{2\pi}{9} + \frac{2\pi}{3}k$

(b)  $\sin^2(x) - \sin(x) = 0$

Factoring,  $\sin x(\sin x - 1) = 0$ , so  $\sin x = 0$  or  $\sin x = 1$

Therefore,  $x = 0 + 2\pi k$  or  $x = \pi + 2\pi k$  or  $x = \frac{\pi}{2} + 2\pi k$

7. Find the values of  $x$ ,  $y$  and  $z$  in the triangle shown below:




First,  $z = 180 - 23 - 90 = 67^\circ$

Next,  $\sin 23^\circ = \frac{y}{6}$ , so  $y = 6 \sin 23^\circ \approx 2.3444$

Similarly,  $\cos 23^\circ = \frac{x}{6}$ , so  $x = 6 \cos 23^\circ \approx 5.5230$  (or we could use the Pythagorean Theorem to find the third side).

8. A function  $f$  is graphed below. Find the following:



(a)  $f(-5)$ ,  $f(-3)$ , and  $f(4)$

From the graph we see  $f(-5) = 0$ ,  $f(-3) = -2$ , and  $f(4) = 2$

(b) find the domain and range of  $f$

Domain:  $(-\infty, 2) \cup (2, \infty)$

Range:  $(-3, \infty)$

(c) find the intervals where  $f$  is decreasing

Decreasing:  $(-\infty, -5) \cup (2, 4)$

(d) find  $\lim_{x \rightarrow 4} f(x)$

$$\lim_{x \rightarrow 4} f(x) = -3$$

(e) find  $\lim_{x \rightarrow 2^-} f(x)$  and  $\lim_{x \rightarrow 2^+} f(x)$

$$\lim_{x \rightarrow 2^-} f(x) = \infty \text{ and } \lim_{x \rightarrow 2^+} f(x) = 1$$

(f) find  $\lim_{x \rightarrow -\infty} f(x)$  and  $\lim_{x \rightarrow \infty} f(x)$

$$\lim_{x \rightarrow -\infty} f(x) = 1 \text{ and } \lim_{x \rightarrow \infty} f(x) = 3$$

(g) find the points where  $f(x)$  is discontinuous, and classify each point of discontinuity.

Points of discontinuity:

$x = -3$  (jump discontinuity)

$x = 2$  (infinite discontinuity)

$x = 4$  (removable discontinuity)

9. Find the following limits:

(a)  $\lim_{x \rightarrow 2} \frac{3x + 7}{\sqrt{5x - 1}} = \frac{3(2) + 7}{\sqrt{5(2) - 1}} = \frac{13}{\sqrt{9}} = \frac{13}{3}$

(b)  $\lim_{x \rightarrow \frac{3}{2}} \frac{2x^2 + x - 6}{4x^2 - 4x - 3} = \lim_{x \rightarrow \frac{3}{2}} \frac{(2x - 3)(x + 2)}{(2x - 3)(2x + 1)} = \lim_{x \rightarrow \frac{3}{2}} \frac{x + 2}{2x + 1} = \frac{\frac{3}{2} + 2}{(2)(\frac{3}{2}) + 1} = \frac{\frac{7}{2}}{4} = \frac{7}{8}$

(c)  $\lim_{x \rightarrow 2} \frac{x^4 - 16}{x^2 - x - 2} = \lim_{x \rightarrow 2} \frac{(x^2 + 4)(x^2 - 4)}{(x - 2)(x + 1)} = \lim_{x \rightarrow 2} \frac{(x^2 + 4)(x - 2)(x + 2)}{(x - 2)(x + 1)} = \lim_{x \rightarrow 2} \frac{(x^2 + 4)(x + 2)}{x + 1}$   
 $= \frac{(2^2 + 4)(2 + 2)}{2 + 1} = \frac{(8)(4)}{3} = \frac{32}{3}$

(d)  $\lim_{x \rightarrow -2^+} \sqrt{x + 2}$

Notice that  $\sqrt{x + 2}$  is defined for  $x \geq -2$ . Therefore,  $\lim_{x \rightarrow -2^+} \sqrt{x + 2} = \sqrt{-2 + 2} = 0$

(e)  $\lim_{x \rightarrow 3^+} \frac{4}{\sqrt{x - 3}}$

Notice that for  $x > 3$ ,  $\sqrt{x - 3} > 0$ . Therefore,  $\lim_{x \rightarrow 3^+} \frac{4}{\sqrt{x - 3}} = \infty$

(f)  $\lim_{x \rightarrow \infty} \frac{(3x - 5)(2x - 3)}{(2x + 1)(3x - 2)}$

$$\lim_{x \rightarrow \infty} \frac{(3x - 5)(2x - 3)}{(2x + 1)(3x - 2)} = \lim_{x \rightarrow \infty} \frac{6x^2 - 19x + 15}{6x^2 - x - 2} = \lim_{x \rightarrow \infty} \frac{x^2(6 - \frac{19}{x} + \frac{15}{x^2})}{x^2(6 - \frac{1}{x} - \frac{2}{x^2})} = \lim_{x \rightarrow \infty} \frac{(6 - \frac{19}{x} + \frac{15}{x^2})}{(6 - \frac{1}{x} - \frac{2}{x^2})} = \frac{6}{6} = 1$$


(g)  $\lim_{x \rightarrow \infty} \frac{(3x - 5)(2x - 3)}{(2x + 1)}$

$$\lim_{x \rightarrow \infty} \frac{(3x - 5)(2x - 3)}{(2x + 1)} = \lim_{x \rightarrow \infty} \frac{6x^2 - 19x + 15}{(2x + 1)} = \lim_{x \rightarrow \infty} \frac{6x - 19 + \frac{15}{x}}{(2 + \frac{1}{x})} = \lim_{x \rightarrow \infty} \frac{6x - 19}{(2)} = \lim_{x \rightarrow \infty} 3x - \frac{19}{2} = \infty$$

10. Given the function

$$f(x) = \begin{cases} x^2 & \text{if } x < 1 \\ 2 & \text{if } x = 1 \\ 4 - x^2 & \text{if } x > 1 \end{cases}$$

(a) Graph  $f(x)$ .



(b) Find  $\lim_{x \rightarrow 1^-} f(x)$ ,  $\lim_{x \rightarrow 1^+} f(x)$ , and  $\lim_{x \rightarrow 1} f(x)$

$$\lim_{x \rightarrow 1^-} f(x) = 1$$

$$\lim_{x \rightarrow 1^+} f(x) = 3$$

$$\lim_{x \rightarrow 1} f(x) \text{ does not exist}$$

(c) Is  $f(x)$  continuous at  $x = 1$ ? Justify your answer.

No. Since  $\lim_{x \rightarrow 1} f(x)$  does not exist,  $f(x)$  is **not** continuous at  $x = 1$ .

11. Given that  $f(x) = x^3 + 5$ ,  $\lim_{x \rightarrow 2} f(x) = 13$ , and  $\epsilon = .01$ , find the largest  $\delta$  such that if  $0 < |x - 2| < \delta$ , then  $|f(x) - 13| < \epsilon$ .

If  $|f(x) - 13| < \epsilon$ , then  $|x^3 + 5 - 13| < \epsilon$ , or  $|x^3 - 8| < .01$

That is,  $-.01 < x^3 - 8 < .01$ , or  $7.99 < x^3 < 8.01$ . Thus  $\sqrt[3]{7.99} < x < \sqrt[3]{8.01}$

Notice that  $2 - \sqrt[3]{7.99} \approx -.000833681$  and  $\sqrt[3]{8.01} - 2 \approx .000832986$

Then the largest  $\delta$  that works is  $\delta = \sqrt[3]{8.01} - 2$

12. Use the formal definition of a limit to prove that  $\lim_{x \rightarrow 6} 5x - 21 = 9$ .

Let  $\epsilon > 0$  be given and suppose that  $|f(x) - 9| < \epsilon$ . Then  $|5x - 21 - 9| = |5x - 30| < \epsilon$ .

But then  $5|x - 6| < \epsilon$ , so  $|x - 6| < \frac{\epsilon}{5}$ .

Therefore, let  $\delta \leq \frac{\epsilon}{5}$ , and suppose  $|x - 6| < \delta$ .

Then  $5|x - 6| < 5\delta \leq \epsilon$ .

Therefore  $|5x - 30| = |5x - 21 - 9| < \epsilon$ , or  $|f(x) - 9| < \epsilon$ .

Thus  $\lim_{x \rightarrow 6} 5x - 9 = 21$

13. Let  $f(x) = \frac{x^2 - x - 2}{x^2 - 2x}$ .

(a) Find the values of  $x$  at which  $f$  is discontinuous.

$$\text{Factoring, } f(x) = \frac{x^2 - x - 2}{x^2 - 2x} = \frac{(x - 2)(x + 1)}{x(x - 2)}$$

Therefore, we can see that  $f(x)$  is discontinuous at  $x = 0$  and  $x = 2$

(b) Find all vertical and horizontal asymptotes of  $f$ .

Since we can cancel the two  $(x - 2)$  terms, there is **not** a vertical asymptote when  $x = 2$

The only vertical asymptote is at  $x = 0$ .

To find the horizontal asymptote, we compute  $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{x^2 - x - 2}{x^2 - 2x} = \lim_{x \rightarrow \infty} \frac{1 - \frac{1}{x} - \frac{2}{x^2}}{1 - \frac{2}{x}} = 1$ .

Thus,  $y = 1$  is the horizontal asymptote of  $f(x)$ .

14. Find the  $x$  values at which  $f(x) = \frac{\sqrt{9 - x^2}}{x^4 - 16}$  is continuous.

First, notice that that  $x^4 - 16 = (x^2 + 4)(x^2 - 4) = (x^2 + 4)(x - 2)(x + 2)$ , so  $f(x)$  is undefined when  $x = 2$  and  $x = -2$ .

Also, for  $f(x)$  to be defined, we must have  $9 - x^2 \geq 0$ , or  $x^2 \leq 9$ . Thus  $-3 \leq x \leq 3$ .

Therefore,  $f(x)$  is continuous on the intervals:  $[-3, -2) \cup (-2, 2) \cup (2, 3]$

15. Use the Intermediate Value Theorem to show  $x^5 - 3x^4 - 2x^3 - x + 1 = 0$  has a solution between 0 and 1.

Let  $f(x) = x^5 - 3x^4 - 2x^3 - x + 1$ . Notice that  $f$  is continuous since it is a polynomial. Also,  $f(0) = 1$  and  $f(1) = -4$ .

Thus, by the IVT, for every  $-4 < w < 1$ , there is a  $c$  satisfying  $0 \leq c \leq 1$  with  $f(c) = w$ . In particular,  $f(c) = 0$  for some  $c$  between zero and 1.