**JASPERSE CHEM 350** TEST 1

VERSION 3 Organic Chemistry I - Jasperse

Intro and Review

Structure and Properties of Organic Molecules

Structure, Nomenclature, and Conformation/Stereochemistry of Alkanes

1. (12 points) Give the relationship between the following pairs of structures. The possible relationships are the following:

same compound stereo isomers

structural isomers resonance structures not isomers (different molecular formula)

a.

Structural. 1,2-dibromo vs 1,3-dibromo.

1. Resonance: No atoms can move! 2. Stereo: same condensed formula 3. Structural: different

condensed formula

H 🚱


ΗН BrBr

Same. **Bond rotation around** single bonds is allowed.

Structural CH3CHCHBr vs CH2CHCH2Br

Stereo Trans-cis. Double bond can't rotate.

CH CH CH CH,



Same. Both are 4-methylnonane

Resonance. Electrons and charge is repositioned, but no atoms moved.

(8 points) Draw line-angle structures and names for 4 of the 5 structural isomers of C<sub>6</sub>H<sub>14</sub>.

3-methylpentane

Alkane Acyclic: C<sub>n</sub>H<sub>2n+2</sub> Alkane Cyclic C<sub>n</sub>H<sub>2n</sub>

Beware of drawing same

2,3-dimethylbutane





- 3. (10 Points)
- a. For the above structure, what is the hybridization and approximate bond angles (109, 120, or 180) about:

- b. In the above structure, N-1 is actually found to have 120° bond angles (This may seem unexpected to you at this point, but we'll learn why later in the course.) What must be the hybridization of the nitrogen?
- sp2.) Hybridization, bond angle, and electron geometry are all interlocked.

  To know any one of them is to know the others.
- 4. (2 Points) Bond rotation around C6-C7 in the above structure has a 7 kcal/mol barrier, while rotation around the C4-C5 bond has a 70 kcal/mol barrier. Explain very briefly why it is so much harder to rotate the latter bond?


Single bond versus double bond.

A double bond has overlapping p-orbitals.

To rotate a double bond, the p-p overlap would be lost. The full pi-bond would need to break.


By contrast, no bonds are broken when you rotate around a single bond.


5. (4 points) For each of the pairs listed, circle the one with the higher boiling point.




- 6. (6 points) Write a Lewis structure and assign any non-zero formal charges.
- 1. Want normal bonding for all: in absence of metal ions
- 2. C=O (or C=C in other cases) may help
- 3. If you have any formal charges (1997), they must sur to net charge of molecule (zero, in this case)
- 4. Organization must match condensed formula sequence










2 Factors:

- More bonds (priority)
- 2. Electronegativity (if bonds are equal)

b. For the two resonance structures shown below, circle the resonance structure that would make the greater contribution to the resonance hybrid.



8. (6 points) Rank the acidity of the following molecules, 1 being most acidic, 4 being least acidic. Hint: draw the anions!



NH<sub>3</sub> 4 CH<sub>3</sub>CH<sub>2</sub>OH <sup>3</sup>

CH<sub>3</sub>CO<sub>2</sub>H

HCI


3 Acidity Factors:


1. charge (not relevant here)


2. Ened 3. Resonance

resonance most stable


9. (6 points) Draw a line-angle picture for all of the atoms in the molecule CH<sub>3</sub>CH<sub>2</sub>COCHClCH<sub>3</sub>, including the hydrogens. Use the hash-wedge convention to indicate atoms that are not in the plane of the paper.








CI could equally well be drawn in the hashed spot










B: By taking on chair conformation, there is zero angle strain, and zero torsional (no eclipsing)

C: If it has ideal angles, then some eclipsing and torsional strain destabilizes it




11. (6 points) Which of the following are capable of cis-trans stereoisomerism? (Yes/No).







b. Expentene (name means a double bond is between carbons 3 and 4)




12. (9 points) Identify the functional groups in the following molecules. (Do not include "alkane", since that is not "functional". And do not specify "cyclic".)










- Longest chain
- 2. Alphabetize substituents
- 3. Number from end near substituent




- 1. cis/trans for di-subbed rings
- 2. Alphabetize substituents
- 3. Numbering
- 4. Know isopropyl and t-butyl


4-ethyl-3,6-dimethyloctane

cis-1-methyl-3-propylcyclobutane or

cis-3-methyl-1-propylcyclobutane


- 14. (8 points) a. Draw Newman projections for the totally eclipsed, the gauch, and the anti conformations of 2,5-dimethylhexane, relative to the C3-C4 bond. You may abbreviate the isopropyl groups attached to C3 and C4 as "i-Pr" for convenience.
- b. Explain very briefly why the rotation barrier around the C3-C4 bond of 2,5-dimethylhexane is greater than the rotation barrier in butane.





b.) Circle the more stable conformation.

c) Would trans-3-methyl-1-isopropylcyclohexane be more stable or less stable than the cis isomer?



- 1. Make sure you've really drawn "flipped" chairs
- 2. What's "ax" in one chair flip is "eq" in the other.
- 3. Process cis-trans
- 4. Draw in H's on substituted carbons (easier to see ax/eq).