


Ch. 11 Liquids, Solids, and Materials

Ch. 15 The Chemistry of Solutes and Solutions

Ch. 13 Chemical Kinetics

Formulas for First Order Reactions: $kt = \ln ([A_0]/[A_t])$ $kt_{1/2} = 0.693$

- Which of the following would have the highest molar heat of vaporization?
a. I_2 b. Br_2 c. Cl_2 d. F_2
- Which of the following would have the highest vapor pressure at $25^\circ C$?
a. C_4H_{10} b. $NaCl$ c. $C_6H_{12}O_6$ d. $C_4H_9NH_2$
- Arrange CH_3OH , NaF , and CO_2 in order of increasing boiling point.
a. $CH_3OH < CO_2 < NaF$
b. $CO_2 < NaF < CH_3OH$
c. $CO_2 < CH_3OH < NaF$
d. $NaF < CO_2 < CH_3OH$
e. none of the above
- Region "c" on the heating curve shown (Temperature versus heat, "q") corresponds to:
a. a pure gas increasing in temperature
b. a liquid increasing in temperature
c. a solid increasing in temperature
d. a solid melting
e. a liquid boiling




- Which of the following would have the highest boiling point?
a. $CH_3CH_2CH_2OH$ b. $CH_3CH_2OCH_3$ c. CH_3CH_2OH d. $CH_3CH_2CH_2CH_3$

6. The H_{fusion} for water is 6.0 kJ/mol, and the specific heat for liquid water is 4.18 J/g-K. How many kJ of heat would it take to convert 36 g of solid ice (18 g/mol) from 0°C to liquid water at 53°C?

a. 8.0 kJ b. 12.0 kJ c. 20.0 kJ d. 7987 kJ

7. In which phase does the substance whose phase diagram is shown below exist at 0°C and atmospheric pressure?

a. gas b. liquid c. solid d. supercritical fluid



8. Which of the following would have the greatest surface tension at 25°C?

a. CH_4 b. CH_3F c. CH_3OH d. CO

9. Which of the following shows a relatively high boiling temperature due to hydrogen bonding?

a. CH_3OH b. CH_3SH c. CH_3OCH_3 d. SnH_4

10. Which of the following substances has London dispersion forces as its only intermolecular force?

a. HCN b. CH_4 c. NH_3 d. H_2S

11. Which is a gas at room temperature? (You may apply memory as well as principle to answer this question!)

a. Na_2S b. NO_2 c. H_2O d. Fe

12. Which is a brittle, high-melting solid but dissolves in water?

- a. I_2 b. K_2CO_3 c. $C_{12}H_{26}$ d. Al

13. Which of the following is not characteristic of a metallic solid?

- a. High electrical conductivity
- b. High thermal conductivity
- c. Good water solubility
- d. Variable but often high melting point
- e. Variable hardness, but malleable as opposed to brittle


14. Which of the following statements is true?

- a. All of the molecules in the liquid state have the same energy
- b. When evaporation occurs the average kinetic energy of the molecules remaining in the liquid state is lower than that of the molecules that left, resulting in cooling of the liquid
- c. The vapor pressure of a liquid decreases as the temperature increases
- d. The rate of evaporation is faster for substances with lower vapor pressures than for substances with higher vapor pressures

15. In a liquid, the attractive intermolecular forces are:

- a. weaker than in a substance that is a gas at the same temperature
- b. always insignificant and unable to keep molecules close together
- c. so strong that molecules are locked close together and are unable to move
- d. strong enough to hold molecules relatively close but not strong enough to keep molecules from moving past each other

16. Which of the following statements is false for the vapor pressure/temperature diagram shown:?



- a. the vapor pressure for C at 60° is about 0.4 atm
- b. substance D has the weakest binding forces
- c. the normal boiling point for A is about 58°
- d. to achieve a vapor pressure of 0.4 atm, substance D must be heated to about 100°C

17. Which of the following statements is false?
- diamond is higher melting than $\text{CH}_3\text{CH}_2\text{OH}$ (alcohol)
 - solid glucose is less dense than melted, liquid glucose
 - CH_2Br_2 is more volatile than CBr_4 at room temperature
 - evaporation of freon-12 absorbs heat from the surroundings
18. Potassium hydroxide dissolves readily in water due to
- strong solute-solute interactions
 - strong solvent-solvent interactions
 - strong solute-solvent interactions
 - weak solute-solvent interactions
19. Which of the following substances would be the most soluble in water?
- Ar
 - $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$
 - NaCl
 - CH_4
20. Which of the following effects would not result when CaCl_2 was added to water?
- the melting point would decrease
 - the boiling point would decrease
 - the vapor pressure would decrease
 - the electrical conductivity of the solution would increase
21. Which relationship is false for solubility in water?
- $\text{C}_5\text{H}_{11}\text{OH} > \text{C}_{11}\text{H}_{23}\text{OH}$
 - $\text{C}_5\text{H}_{11}\text{OH} > \text{C}_5\text{H}_{12}$
 - $\text{CCl}_4 > \text{CaCl}_2$
 - $\text{CH}_3\text{OCH}_3 > \text{CH}_3\text{CCl}_3$
22. Which of the following statements is true?
- In a saturated solution at equilibrium, the rate at which solid material is dissolving differs from the rate at which solid material is reforming
 - For solids that dissolve in water, the primary reason is because dissolving results in increasing disorder
 - Any solid that dissolves in water does so in an exothermic way
 - The solubility of a solid always decreases at higher temperature
23. Which of the following should be least miscible in carbon tetrachloride, CCl_4 ?
- C_6H_{14}
 - CH_3OH
 - Br_2
 - C_3H_8
24. Which one of the following 0.1 M aqueous solutions would have the lowest melting/freezing point?
- $\text{CH}_3\text{CH}_2\text{OH}$
 - AlPO_4
 - NaNO_3
 - CaBr_2

25. Consider the following four solutions, and choose which statement is false :
- 1 L of Pure Water
 - 1 L of water with 0.15 moles of CH_3OH added
 - 1 L of water with 0.15 moles of $\text{CH}_3\text{CH}_2\text{OH}$
 - 1 L of water with 0.15 moles of NaCl added
- The pure water solution will have the highest vapor pressure
 - The solution with 0.15 moles of CH_3OH will have the same vapor pressure as the solution with 0.15 moles of $\text{CH}_3\text{CH}_2\text{OH}$
 - The solution with 0.15 moles of CH_3OH will have higher vapor pressure than the solution with 0.15 moles of NaCl
 - The solution with 0.15 moles of NaCl will have the highest vapor pressure
26. Which of the following statements is false?
- C_6H_{14} has very low solubility in water because it can't hydrogen bond to itself or to water
 - NaCl has poor solubility in CCl_4 because strong solute-solute interactions are replaced by feeble solute-solvent interactions, making things strongly endothermic
 - C_6H_{14} has good solubility in CCl_4 . Neither original nor final intermolecular interactions are very strong.
 - CH_3OCH_3 has very low solubility in water because it can't hydrogen bond to itself or to water
27. If the reaction $2\text{A} + 3\text{D} \rightarrow \text{products}$ is first-order in A and second-order in D, then the rate law will have the form, rate =
- $k[\text{A}][\text{D}]$
 - $k[\text{A}]^2[\text{D}]^3$
 - $k[\text{A}][\text{D}]^2$
 - $k[\text{A}]^2[\text{D}]$
 - $k[\text{A}]^2[\text{D}]^2$
28. A reaction has rate law, rate = $k[\text{A}]^2$. What are the units of k?
- M/s
 - $\text{M}^{-1} \text{s}^{-1}$
 - 1/s
 - 1/M
 - s/M^2
29. Consider the reaction $\text{A} + \text{B} \rightarrow 4\text{C}$, if the rate of disappearance of A is 0.16 mol/min, what is the rate of formation of C?
- 0.04 mol/min
 - 0.16 mol/min
 - 0.32 mol/min
 - 0.64 mol/min
 - none of the above
30. What is the rate law for the reaction $\text{A} + 3\text{B} \rightarrow \text{products}$
- | Initial [A] | Initial [B] | rate |
|-------------|-------------|------|
| 0.273 | 0.763 | 3.0 |
| 0.273 | 1.526 | 3.0 |
| 0.819 | 0.763 | 27.0 |
- rate = $k[\text{A}][\text{B}]$
 - rate = $k[\text{A}]$
 - rate = $k[\text{A}]^2$
 - rate = $k[\text{A}]^3$
 - none of the above

31. What is the rate constant k (ignore units) for the reaction shown, if the reaction is first order in both A and B.



Initial [A]	Initial [B]	rate
0.23	0.17	0.33

- a. 8.4
- b. 5.6
- c. 0.67
- d. 0.18

32. What is the rate law for the reaction $2A + 5B \rightarrow$ products

Initial [A]	Initial [B]	rate
0.421	0.234	6.4×10^4
0.842	0.234	1.3×10^5
0.421	0.468	2.6×10^5

- a. $\text{rate} = k[A][B]$
- b. $\text{rate} = k[B]$
- c. $\text{rate} = k[A][B]^2$
- d. $\text{rate} = k[A][B]^3$
- e. none of the above

33. If the rate law for a reaction is $\text{rate} = k[A]^2[B]$, what is the effect on the overall rate of doubling the concentration of both A and B?

- a. rate increases by 2
- b. rate increases by 4
- c. rate increases by 8
- d. rate increases by 16
- e. none of the above

34. $A \rightarrow B$ is a first order reaction. If $k = 6.30 \times 10^{-4} \text{ s}^{-1}$, and the initial $[A] = 0.100 \text{ M}$, what is $[A]$ after 1000 s?


- a. 0.0533
- b. 0.0234
- c. 0.188
- d. 0.427
- e. 0.000100

35. $A \rightarrow B$ is a first order reaction. What is the rate constant for the reaction (in s^{-1}).

time (sec)	[A] (M)
0.0	1.60
5.0	0.80
10.0	0.40
15.0	0.20
20.0	0.10

- a) 0.013
- b) 0.030
- c) 0.14
- d) 3.0

36. For the reaction diagram shown, which of the following statements is true?



- Line W represents the ΔH for the forward reaction; point B represents the transition state
- Line W represents the activation energy for the forward reaction; point B represents the transition state
- Line Y represents the activation energy for the forward reaction; point C represents the transition state
- Line X represents the ΔH for the forward reaction; point B represents the transition state

37. Given the mechanism shown, what would be the rate law?



- rate = $k[\text{NO}]^2[\text{Br}_2]$
- rate = $k[\text{N}_2\text{O}_2]^2[\text{Br}_2]$
- rate = $k[\text{NO}]^2[\text{N}_2\text{O}_2][\text{Br}_2]$
- rate = $k[\text{NO}][\text{Br}_2]$
- rate = $k[\text{NO}]$

38. A catalyst increases the reaction rate by

- always reducing the number of elementary steps in the mechanism
- always making the overall transition state higher in energy
- changing the mechanism to lower the overall activation energy barrier

39. In any multistep reaction mechanism, the rate of the overall reaction is determined by the rate of the _____ step in the mechanism.

- first
- last
- slowest
- fastest

40. Which of the following statements is true?

- a. the activation energy always increases as temperature rises
- b. the activation energy always decreases as temperature rises
- c. the rate constant always decreases as temperature rises
- d. the rate constant always increases as temperature rises
- e. the rate constant always increases as the activation energy increases

41. Which of the following statements is false regarding collision theory?

- a. As temperature rises, a higher number of bimolecular collisions result in successful reaction
- b. As the concentration of either chemical increases, the bimolecular collision frequency increases
- c. Not all bimolecular collisions result in successful reactions
- d. Elementary steps are routine that are either termolecular (three molecules colliding at once) or tetramolecular (four molecules colliding at once)

Jasperse
Chem 160
Test1
Version 1
Answers

- | | |
|-------|-------|
| 1. A | 22. B |
| 2. A | 23. B |
| 3. C | 24. D |
| 4. B | 25. D |
| 5. A | 26. D |
| 6. C | 27. C |
| 7. C | 28. B |
| 8. C | 29. D |
| 9. A | 30. C |
| 10. B | 31. A |
| 11. B | 32. C |
| 12. B | 33. C |
| 13. C | 34. A |
| 14. B | 35. C |
| 15. D | 36. B |
| 16. B | 37. A |
| 17. B | 38. C |
| 18. C | 39. C |
| 19. C | 40. D |
| 20. B | 41. D |
| 21. C | |