

Chapter 20 Nuclear Chemistry Math Summary

Particles Involved in Nuclear Reactions, either as Nucleons, Emitted particles or Particles that React with a Nucleus and Induce a Decay

(Memorize these for Test)

-the first three, alpha, beta, and positrons are the crucial ones for balancing radioactive nuclear decay reactions

$\frac{4}{2}\text{He}$ α -particle (alpha)	$\frac{0}{0}\gamma$ gamma
$\frac{0}{-1}\text{e}$ β -particle (beta), electron	$\frac{1}{0}\text{n}$ neutron
$\frac{0}{+1}\text{e}$ positron	$\frac{1}{1}\text{H}$ proton

Radioactive Decay Math

$t = (t_{1/2}/0.693) \ln (A_0/A_t)$ When solving for time, given half life and quantities of material

$\ln (A_0/A_t) = 0.693 (t /t_{1/2})$ When solving for the amount of material left after a given time, given the half life

Handling “ $\ln y = x$ ” on calculator, when you know “ x ” but want to solve for “ y ”: enter “ x ”, then hit your “ e^x ” button.

Mass Defect/Binding Energy Math

Proton mass: 1.00783

Neutron mass: 1.00867

$$E = \Delta mc^2$$

$$\Delta m = (\text{sum mass of protons plus neutrons}) - \text{actual mass}$$

- The binding energy will depend on the Δm difference between the summed weight of the protons and neutrons minus the actual mass of the nucleus.
- Δm in terms of kilograms (you’ll normally need to convert from grams to kg)
- The energy answer from the formula comes out in terms of Joules, not kJ; you’ll routinely need to convert from J to kJ to fit the answers