


Exam 3 – Electrodynamics

April 5, 2007

This is a closed book examination. There is extra scratch paper available. Please explain your answers. Your explanation is worth 3/4 of the points on multiple-choice questions.

- 1) [4 PTS] What is the voltage at $t = 0.3$ s for a sinusoidal voltage source with $V_{\text{rms}} = 4.0$ V and $f = \frac{1}{4}$ Hz? Note: $V(t=0\text{s}) = 0$ V. WRITE OUT THE EQUATION.
- $0.30 \pm .02$ V
 - $0.42 \pm .02$ V
 - $1.81 \pm .02$ V
 - $2.57 \pm .02$ V
 - $5.27 \pm .02$ V
- 2) [4 PTS] A positively charged object is placed in the center of a room with a uniform magnetic field. If the magnetic field is pointing east the object will
- move east.
 - move west.
 - move north.
 - move south.
 - not move.
- 3) [4 PTS] A 1.5 V battery is connected to a transformer on a side with 1000 windings. The other side has only 10 windings so the output voltage is
- 150 V
 - 15 V
 - 0 V
 - 0.15 V
 - 0.015 V
- 4) [4 PTS] A wire bent into a “C” shape moves at a constant speed perpendicular to (and away from) a long straight current-carrying wire.
- The induced current in the loop will progress clockwise.
 - The induced current in the loop will progress counterclockwise.
 - There will be no induced current.
 - The induced current decrease as the loop moves away
 - None of the above




- 5) [4 PTS] When the effective (rms) voltage and current in an ac circuit are in-phase, we know
- the total reactance is zero
 - the capacitive reactance is zero
 - the inductive reactance is zero
 - the resistance is zero
 - the impedance is zero
 - both (b) and (d)

- 6) [4 PTS] An inductor and resistor are connected in series to an AC voltage source. If you double the frequency of the voltage the effect on the inductor is to
- double its reactance
 - increase its reactance by a factor of four
 - leave its reactance unchanged
 - halve its reactance
 - decrease its reactance by a factor of four


- 7) [4 PTS] If you double the current passing through an inductor, the energy stored in the inductor
- increases $E_i = \frac{1}{4}E_f$
 - increases $E_i = \frac{1}{2}E_f$
 - does not change.
 - decreases $E_i = 2E_f$
 - decreases $E_i = 4E_f$

- 8) [4 PT] Two very long wires, 45 cm apart, are hung parallel to each other. Current flows down each wire in the same direction. Wire C has a current of 2 Amps and wire B has a current of 4 Amps.

- The two wires are attracted $F_C = \frac{1}{4}F_B$
- The two wires are attracted $F_C = \frac{1}{2}F_B$
- The two wires are attracted $F_C = F_B$
- The two wires are repelled $F_C = F_B$
- The two wires are repelled $F_C = 2F_B$
- The two wires are repelled $F_C = 4F_B$



- 9) [10 PT] A square loop of wire is placed in a uniform magnetic field, $B=1.5$ T. The loop has sides that are 2 cm and is rotated at 2 Hz perpendicular to the magnetic field. Two leads connect to the loop.
- What is the maximum magnetic flux in the loop?
 - Graph the voltage generated across the leads as a function of time.
 - List (at least) two ways you could increase the voltage across the leads.



- 10) [10 PT] A solid silver wire of diameter 2 mm carries a current of 10 mA. The current density is constant and uniform. Using Ampere's Law graph the magnetic field inside and outside the wire.

Useful mathematical relationships:

$$\sin^2(\theta) + \cos^2(\theta) = 1 \quad \text{and} \quad \sin(2\theta) = 2 \sin(\theta) \cos(\theta)$$

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta) = 2 \cos^2(\theta) - 1 = 1 - 2 \sin^2(\theta)$$

$$\int_b^c \frac{a}{x} = a \ln x \Big|_b^c = a [\ln c - \ln b] = a \ln\left(\frac{c}{b}\right)$$

